

Parametrization of the isospectral set for the vector-valued Sturm-Liouville problem

Dmitry Chelkak* and Evgeny Korotyaev†

February 2, 2008

Abstract

We obtain a parametrization of the isospectral set of matrix-valued potentials for the vector-valued Sturm-Liouville problem on a finite interval.

AMS Classification: 34A55, 34B24.

Keywords: Sturm-Liouville problem, matrix potentials, isospectral potentials.

1 Introduction and main results

Consider the inverse problems for the self-adjoint operator H in $L^2(0, 1)^N$, given by

$$H\psi = -\psi'' + V(x)\psi = \lambda\psi, \quad \psi(0) = \psi(1) = 0, \quad (1.1)$$

where $V = V^* \in L^1(0, 1)$ is some $N \times N$ matrix-valued potential and ψ is a vector-valued function. Let

$$\lambda_1 < \lambda_2 < \dots < \lambda_\alpha < \dots$$

be the eigenvalues of H , where each λ_α , $\alpha \geq 1$, has multiplicity $k_\alpha \in [1, N]$, i.e. k_α is the number of eigenfunctions corresponding to the eigenvalue λ_α .

In the scalar case this problem is well-known, including the complete characterization of the set of spectral data (eigenvalues and norming constants) that correspond to various classes of potentials (see [L], [M], [PT]). On the contrary, in the matrix case only some particular results are known. Thus, it is known that the matrix-valued Weyl-Titchmarsh function (see (1.5)) uniquely determines the potential V (see [Mal] or [Yu], where this Borg-type uniqueness theorem was proved in two different ways). Jodeit and Levitan [JL1] and Chern [Ch] have constructed some isospectral sets of potentials, which not only have the same spectrum as V but also the same initial data (boundary values) of eigenfunctions.

*Dept. of Math. Analysis, Math. Mech. Faculty, St.Petersburg State University. Universitetskij pr. 28, Staryj Petergof, 198504 St.Petersburg, Russia, e-mail: delta4@math.spbu.ru

†Institut für Mathematik, Humboldt Universität zu Berlin, Rudower Chaussee 25, 12489, Berlin, Germany, e-mail: evgeny@math.hu-berlin.de

Some other results about the inverse spectral problem for the vector-valued Schrödinger equation were obtained in [CHGL], [CK], [Ca], [ChSh], [JL2], [Sh], [SP].

We denote by $\varphi(x, \lambda)$ the $N \times N$ matrix-valued solution of the equation $-\varphi'' + V\varphi = \lambda\varphi$ such that $\varphi(0, \lambda) = 0$ and $\varphi'(0, \lambda) = I$, where $I = I_N$ is the identity matrix. Introduce the matrices

$$S_\alpha = \int_0^1 (\varphi^* \varphi)(t, \lambda_\alpha) dt = S_\alpha^* > 0, \quad \alpha \geq 1.$$

Definition 1.1 (Spectral data). For each eigenvalue λ_α , $\alpha \geq 1$, we define the subspace

$$\mathcal{E}_\alpha = \text{Ker } \varphi(1, \lambda_\alpha) = \{h \in \mathbb{C}^N : \varphi(1, \lambda_\alpha)h = 0\} \subset \mathbb{C}^N, \quad \dim \mathcal{E}_\alpha = k_\alpha, \quad (1.2)$$

the orthogonal projector $P_\alpha : \mathbb{C}^N \rightarrow \mathcal{E}_\alpha$ and the positive self-adjoint operator $g_\alpha : \mathcal{E}_\alpha \rightarrow \mathcal{E}_\alpha$ given by

$$g_\alpha = G_\alpha|_{\mathcal{E}_\alpha}, \quad \text{where } G_\alpha = P_\alpha S_\alpha P_\alpha.$$

Remark. Each solution of the equation $-\psi'' + V\psi = \lambda\psi$ under the condition $\psi(0) = 0$ has the form $\varphi(x, \lambda)h$ for some $h \in \mathbb{C}^N$. Then, each eigenfunction ψ_α such that $H\psi_\alpha = \lambda_\alpha\psi_\alpha$ has the form $\psi_\alpha(x) = \varphi(x, \lambda_\alpha)h$ for some $h \in \mathcal{E}_\alpha$. In particular, we get $\dim \mathcal{E}_\alpha = k_\alpha$. Moreover, the following identity is fulfilled:

$$\int_0^1 \|\varphi(x, \lambda_\alpha)h\|^2 dx = \langle h, g_\alpha h \rangle, \quad h \in \mathcal{E}_\alpha,$$

where $\langle u, v \rangle = u^*v$ is the scalar product of two vectors and $\|u\|^2 = \langle u, u \rangle$.

Below we will sometimes write $\varphi(x, \lambda, V), \lambda_\alpha(V), \dots$, instead of $\varphi(x, \lambda), \lambda_\alpha, \dots$, when several potentials are being dealt with.

Let $\tilde{\lambda}_\alpha = \lambda_\alpha(\tilde{V})$, $\tilde{\varphi}(x, \lambda) = \varphi(x, \lambda, \tilde{V})$ and so on. For $V = V^* \in L^1(0, 1)$ we introduce the isospectral set of potentials by

$$\text{Iso}(V) = \left\{ \tilde{V} = \tilde{V}^* \in L^1(0, 1) : \tilde{\lambda}_\alpha = \lambda_\alpha, \tilde{k}_\alpha = k_\alpha \text{ for all } \alpha \geq 1 \right\}. \quad (1.3)$$

Our goal is to show that the spectral data $\{(\mathcal{E}_\alpha, g_\alpha)\}_{\alpha \geq 1}$ give the "proper" parametrization of the set $\text{Iso}(V)$. For the sake of the reader, we start with the simple result that the parameters $\{(\lambda_\alpha, \mathcal{E}_\alpha, g_\alpha)\}_{\alpha \geq 1}$ determine the potential uniquely (see Theorem 1.2). In fact, these data are closely related to the residues of the Weyl-Titchmarsh function (see Proposition 1.6). Our main result is Theorem 1.3 which shows that $\{(\mathcal{E}_\alpha, g_\alpha)\}_{\alpha \geq 1}$ are **free parameters**. Namely, we prove that each $\mathcal{E}_\alpha$ and g_α can be changed in an almost arbitrary way, when all other parameters $\{(\mathcal{E}_\beta, g_\beta)\}_{\beta \neq \alpha}$ and the spectrum are fixed.

Theorem 1.2 (Uniqueness). Let $\tilde{V} \in \text{Iso}(V)$ for some $V = V^* \in L^1(0, 1)$ and let $\tilde{\mathcal{E}}_\alpha = \mathcal{E}_\alpha$ and $\tilde{g}_\alpha = g_\alpha$ for all $\alpha \geq 1$. Then $\tilde{V} = V$.

Introduce the "forbidden subspace"

$$\mathcal{F}_\alpha = \mathbb{C}^N \ominus (S_\alpha(\mathcal{E}_\alpha)), \quad \alpha \geq 1, \quad (1.4)$$

where $\mathbb{C}^N \ominus \mathcal{S} = \{v \in \mathbb{C}^N : \langle v, u \rangle = 0 \text{ for all } u \in \mathcal{S}\}$ is the orthogonal subspace to $\mathcal{S} \subset \mathbb{C}^N$. Note that $\dim \mathcal{F}_\alpha = N - k_\alpha$, since $\dim \mathcal{E}_\alpha = k_\alpha$ and $S_\alpha > 0$. We formulate our main result.

Theorem 1.3. *Let $V = V^* \in L^1(0, 1)$ and $\alpha \geq 1$. Then the mapping $\Phi_\alpha : \tilde{V} \mapsto (\tilde{\mathcal{E}}_\alpha, \tilde{g}_\alpha)$ is a bijection between the set of potentials*

$$\left\{ \tilde{V} \in \text{Iso}(V) : \tilde{\mathcal{E}}_\beta = \mathcal{E}_\beta, \tilde{g}_\beta = g_\beta \text{ for all } \beta \neq \alpha \right\}$$

and the following set of pairs $(\tilde{\mathcal{E}}_\alpha, \tilde{g}_\alpha)$:

$$\{(\mathcal{E}, g) : g = g^* > 0 \text{ is an operator in some subspace } \mathcal{E} \subset \mathbb{C}^N$$

$$\text{with } \dim \mathcal{E} = k_\alpha \text{ such that } \mathcal{E} \cap \mathcal{F}_\alpha = \{0\}\}.$$

Remark. i) Substituting $\tilde{V} = V$, we obtain $\mathcal{E}_\alpha \cap \mathcal{F}_\alpha = \{0\}$.

ii) If $\tilde{\mathcal{E}}_\alpha = \mathcal{E}_\alpha$, then there are no restrictions on the changing of the "norming matrix" g_α . This case is similar to the scalar case and [JL1], [Ch].

iii) If we change $\tilde{\mathcal{E}}_\alpha$, then there exists only one restriction $\tilde{\mathcal{E}}_\alpha \cap \mathcal{F}_\alpha = \{0\}$. Such an effect is absent in the scalar case. As far as the authors know, this is the first result in this direction.

iv) Note that we use an explicit procedure (see Theorem 3.4), which is based on the so-called Darboux transform. Therefore, the result of any finite number of such changes can be expressed explicitly in terms of the initial potential.

The next Proposition shows that each "forbidden subspace" $\mathcal{F}_\alpha$, $\alpha \geq 1$, doesn't depend on the "norming matrices" $\{g_\beta\}_{\beta \geq 1}$. Namely, it is uniquely determined by the spectrum and all subspaces $\{\mathcal{E}_\beta\}_{\beta \neq \alpha}$.

Proposition 1.4. *Let $\tilde{V} \in \text{Iso}(V)$ for some $V = V^* \in L^1(0, 1)$. Fix some $\alpha \geq 1$ and let $\tilde{\mathcal{E}}_\beta = \mathcal{E}_\beta$ for all $\beta \neq \alpha$. Then $\tilde{\mathcal{F}}_\alpha = \mathcal{F}_\alpha$ and $\tilde{\mathcal{E}}_\alpha \cap \mathcal{F}_\alpha = \{0\}$.*

In order to illustrate the "forbidden subspaces", we give the following simple example.

Proposition 1.5 (Example). *Let $N = 2$ and $V = V^* \in L^1(0, 1)$ be such that $k_1 = k_2 = 1$ and $k_\alpha = 2$ for all $\alpha \geq 3$. Then $\mathcal{E}_1 \cap \mathcal{E}_2 = \{0\}$, $\mathcal{F}_1 = \mathcal{E}_2$ and $\mathcal{F}_2 = \mathcal{E}_1$.*

Finally, we give the connection between our spectral data and the matrix-valued Weyl-Titchmarsh function $m(\lambda)$ given by

$$m(\lambda) = (\chi' \chi^{-1})(0, \lambda), \quad \lambda \in \mathbb{C}, \quad (1.5)$$

where $\chi(x, \lambda)$ is the matrix-valued solution of the equation $-\chi'' + V\chi = \lambda\chi$ such that $\chi(1, \lambda) = 0$ and $\chi'(1, \lambda) = I$.

Proposition 1.6. *Let $V = V^* \in L^1(0, 1)$. Then the function m is analytic in $\mathbb{C} \setminus \bigcup_{\alpha \geq 1} \{\lambda_\alpha\}$ and satisfies the identity $m(\lambda) = m^*(\bar{\lambda})$. Moreover, each point λ_α , $\alpha \geq 1$, is a simple pole of m and*

$$\left. \text{res}_{\lambda=\lambda_\alpha} m(\lambda) \right|_{\mathcal{E}_\alpha} = -g_\alpha^{-1}, \quad \left. \text{res}_{\lambda=\lambda_\alpha} m(\lambda) \right|_{\mathbb{C}^N \ominus \mathcal{E}_\alpha} = 0, \quad \alpha \geq 1.$$

We describe the plan of the paper. In Sect. 2 we prove some preliminary Lemmas, Theorem 1.2 and Proposition 1.6. In Sect. 3 we prove Theorem 1.3 and Propositions 1.4, 1.5.

2 Preliminaries

Repeating the standard arguments (see [PT], p. 13–15), we obtain the following asymptotics:

$$\varphi(x, z^2) = \frac{\sin zx}{z} \cdot I - \frac{\cos zx}{2z^2} \int_0^x V(t)dt + o\left(\frac{e^{|\operatorname{Im} z|x}}{|z|^2}\right), \quad (2.1)$$

$$\varphi'(x, z^2) = \cos zx \cdot I + \frac{\sin zx}{2z} \int_0^x V(t)dt + o\left(\frac{e^{|\operatorname{Im} z|x}}{|z|}\right), \quad (2.2)$$

as $|z| \rightarrow \infty$ for all $x \in [0, 1]$ and $V \in L^1(0, 1)$. Also, note that

$$\chi(x, \lambda, V) = -\varphi(1-x, \lambda, V^\sharp), \quad \text{where } V^\sharp(t) = V(1-t), \quad t \in [0, 1]. \quad (2.3)$$

Lemma 2.1. *Let $V = V^* \in L^1(0, 1)$. Then*

(i) *for each $(x, \lambda) \in [0, 1] \times \mathbb{C}$ the following identities are fulfilled:*

$$\varphi^*(x, \bar{\lambda})\varphi'(x, \lambda) = (\varphi')^*(x, \bar{\lambda})\varphi(x, \lambda), \quad \chi^*(x, \bar{\lambda})\chi'(x, \lambda) = (\chi')^*(x, \bar{\lambda})\chi(x, \lambda), \quad (2.4)$$

$$\chi^*(0, \bar{\lambda}) = \chi^*(x, \bar{\lambda})\varphi'(x, \lambda) - (\chi')^*(x, \bar{\lambda})\varphi(x, \lambda) = -\varphi(1, \lambda). \quad (2.5)$$

(ii) *for each $\alpha \geq 1$ the following identity is fulfilled:*

$$G_\alpha = P_\alpha [\dot{\varphi}^* \varphi'](1, \lambda_\alpha) P_\alpha. \quad (2.6)$$

Proof. (i) The function $\eta(x) = \chi^*(x, \bar{\lambda})\varphi'(x, \lambda) - (\chi')^*(x, \bar{\lambda})\varphi(x, \lambda)$ satisfies the equation

$$\eta'(x) = \chi^*(x, \bar{\lambda}) ((V(x) - \lambda I) - (V^*(x) - \bar{\lambda} I)^*) \varphi(x, \lambda) = 0, \quad x \in [0, 1].$$

Due to $\eta(0) = \chi^*(0, \bar{\lambda})$ and $\eta(1) = -\varphi(1, \lambda)$, we obtain (2.5). The proof of (2.4) is similar.

(ii) Note that $-\dot{\varphi}'' = (\lambda I - V)\dot{\varphi} + \varphi$. This gives $[\dot{\varphi}^* \varphi' - (\dot{\varphi}')^* \varphi]'(x, \lambda_\alpha) = [\varphi^* \varphi](x, \lambda_\alpha)$, since $\lambda_\alpha \in \mathbb{R}$. Therefore,

$$G_\alpha = P_\alpha [\dot{\varphi}^* \varphi' - (\dot{\varphi}')^* \varphi](1, \lambda_\alpha) P_\alpha = P_\alpha [\dot{\varphi}^* \varphi'](1, \lambda_\alpha) P_\alpha,$$

where we have used $\varphi(1, \lambda_\alpha) P_\alpha = 0$. □

Introduce the matrices

$$Z_\alpha = \dot{\varphi}(1, \lambda_\alpha) P_\alpha + \varphi(1, \lambda_\alpha) P_\alpha^\perp, \quad \text{where } P_\alpha^\perp = I - P_\alpha, \quad \alpha \geq 1. \quad (2.7)$$

Lemma 2.2. *Let $V = V^* \in L^1(0, 1)$. Then (i) $\det Z_\alpha \neq 0$ for all $\alpha \geq 1$.*

(ii) *Each λ_α , $\alpha \geq 1$, is the root of the entire function $\det \varphi(1, \lambda)$ of the multiplicity k_α . The function $\det \varphi(1, \lambda)$ has no other roots. Moreover, the following asymptotics is fulfilled:*

$$\varphi^{-1}(1, \lambda) = ((\lambda - \lambda_\alpha)^{-1} P_\alpha + P_\alpha^\perp)(Z_\alpha^{-1} + O(\lambda - \lambda_\alpha)) \quad \text{as } \lambda \rightarrow \lambda_\alpha. \quad (2.8)$$

(iii) *Let $\xi(\lambda)$, $\lambda \in \mathbb{C}$, be the entire $N \times N$ matrix-valued function such that $\xi(\lambda_\alpha) P_\alpha = 0$ for all $\alpha \geq 1$. Then, $\xi(\lambda) \varphi^{-1}(1, \lambda)$ is the entire matrix-valued function.*

Proof. (i) Suppose that $\dot{\varphi}(1, \lambda_\alpha)P_\alpha h + \varphi(1, \lambda_\alpha)P_\alpha^\perp h = 0$ for some vector $h \in \mathbb{C}^N$. Using (2.6) and (2.4), we obtain

$$\begin{aligned}\langle P_\alpha h, g_\alpha P_\alpha h \rangle &= h^* P_\alpha S_\alpha P_\alpha h = h^* P_\alpha [\dot{\varphi}^* \varphi'](1, \lambda_\alpha) P_\alpha h \\ &= -h^* P_\alpha^\perp [\varphi^* \varphi'](1, \lambda_\alpha) P_\alpha h = -h^* P_\alpha^\perp [(\varphi')^* \varphi](1, \lambda_\alpha) P_\alpha h = 0,\end{aligned}$$

since $\varphi(1, \lambda_\alpha)P_\alpha = 0$. Therefore, $P_\alpha h = 0$ and $\varphi(1, \lambda_\alpha)P_\alpha^\perp h = 0$, i.e. $P_\alpha^\perp h \in \mathcal{E}_\alpha$ and $h = 0$.

(ii) Note that $\det \varphi(1, \lambda) = 0$ if and only if $\varphi(1, \lambda)h = 0$ for some $h \in \mathbb{C}^N$, $h \neq 0$, i.e. if and only if λ is an eigenvalue of the operator H . Let $\lambda - \lambda_\alpha = \mu$. Due to $\varphi(1, \lambda_\alpha)P_\alpha = 0$, we have

$$\varphi(1, \lambda) = (\mu \dot{\varphi}(1, \lambda_\alpha) + O(\mu^2))P_\alpha + (\varphi(1, \lambda_\alpha) + O(\mu))P_\alpha^\perp = (Z_\alpha + O(\mu))(\mu P_\alpha + P_\alpha^\perp)$$

as $\mu \rightarrow 0$. This implies (2.8), since $\det Z_\alpha \neq 0$. Moreover,

$$\det \varphi(1, \lambda) = \det(Z_\alpha + O(\mu)) \det(\mu P_\alpha + P_\alpha^\perp) = (\det Z_\alpha + O(\mu)) \mu^{k_\alpha} \quad \text{as } \mu \rightarrow 0,$$

i.e. the multiplicity of the root λ_α is equal to k_α .

(iii) Note that $\varphi^{-1}(1, \lambda)$ is analytic outside the set $\bigcup_{\alpha \geq 1} \{\lambda_\alpha\}$. Fix some $\alpha \geq 1$. Due to $\xi(\lambda_\alpha)P_\alpha = 0$, we have $\mu^{-1}\xi(\lambda)P_\alpha = O(1)$ as $\mu \rightarrow 0$. It follows from (2.8) that $\xi(\lambda)\varphi^{-1}(1, \lambda)$ is bounded near λ_α for each $\alpha \geq 1$. Therefore, $\xi(\lambda)\varphi^{-1}(1, \lambda)$ is entire. $\square$

Recall that we use the notations $\tilde{\varphi}(x, \lambda) = \varphi(x, \lambda, \tilde{V})$, $\tilde{\mathcal{E}}_\alpha = \mathcal{E}_\alpha(\tilde{V})$ and so on.

Proposition 2.3. (i) Let $\tilde{V} \in \text{Iso}(V)$ for some $V = V^* \in L^1(0, 1)$ and let $\tilde{\mathcal{E}}_\alpha = \mathcal{E}_\alpha$ for all $\alpha \geq 1$. Then

$$\tilde{\varphi}(1, \lambda) = \varphi(1, \lambda) \quad \text{for all } \lambda \in \mathbb{C}.$$

(ii) Let, in addition, $\tilde{g}_\alpha = g_\alpha$ for all $\alpha \geq 1$. Then

$$[\tilde{\varphi}' \tilde{\varphi}^{-1}](1, \lambda) = [\varphi' \varphi^{-1}](1, \lambda) \quad \text{for all } \lambda \in \mathbb{C}.$$

Proof. (i) Due to $\tilde{P}_\alpha = P_\alpha$ and Lemma 2.2 (iii), the function $[\tilde{\varphi} \varphi^{-1}](1, \lambda)$ is entire. Moreover, asymptotics (2.1) gives $[\tilde{\varphi} \varphi^{-1}](1, z^2) = I + O(|z|^{-1})$ as $|z| = \pi(n + \frac{1}{2}) \rightarrow \infty$. Using the Liouville Theorem, we obtain $[\tilde{\varphi} \varphi^{-1}](1, \lambda) = I$ for all $\lambda \in \mathbb{C}$.

(ii) Put

$$f(\lambda) = [(\tilde{\varphi}' \tilde{\varphi}^{-1}) - (\varphi' \varphi^{-1})](1, \lambda) = [(\tilde{\varphi}' - \varphi') \varphi^{-1}](1, \lambda).$$

Firstly, we prove that the function $f(\lambda)$ is entire. Due to Lemma 2.2 (iii), it is sufficient to check that $[\tilde{\varphi}' - \varphi'](1, \lambda_\alpha)P_\alpha = 0$ for all $\alpha \geq 1$. Recall that $Z_\alpha = \dot{\varphi}(1, \lambda_\alpha)P_\alpha + \varphi(1, \lambda_\alpha)P_\alpha^\perp$ and $\det Z_\alpha \neq 0$. Using (2.6) and (2.4), we get

$$Z_\alpha^*[\tilde{\varphi}' - \varphi'](1, \lambda_\alpha)P_\alpha = \tilde{G}_\alpha - G_\alpha + P_\alpha^\perp[(\tilde{\varphi}' - \varphi')^* \varphi](1, \lambda_\alpha)P_\alpha = 0,$$

since $\tilde{G}_\alpha = G_\alpha$ and $\varphi(1, \lambda_\alpha)P_\alpha = 0$. This gives $[\tilde{\varphi}' - \varphi'](1, \lambda_\alpha)P_\alpha = 0$ for all $\alpha \geq 1$.

Secondly, note that asymptotics (2.1), (2.2) yield $f(z^2) = O(1)$ as $|z| = \pi(n + \frac{1}{2}) \rightarrow \infty$, and $f(z^2) \rightarrow 0$ as $z \rightarrow i\infty$. Using the Liouville Theorem, we obtain $f(\lambda) = 0$, $\lambda \in \mathbb{C}$. $\square$

Proof of Theorem 1.2. Recall that $\chi(x, \lambda)$ is the solution of the equation $-\chi'' + V\chi = \lambda\chi$ under the conditions $\chi(1, \lambda) = 0$ and $\chi'(1, \lambda) = I$. Introduce the $2N \times 2N$ matrix

$$K(x, \lambda) = \begin{pmatrix} \tilde{\varphi} & \tilde{\chi} \\ \tilde{\varphi}' & \tilde{\chi}' \end{pmatrix}(x, \lambda) \begin{pmatrix} \varphi & \chi \\ \varphi' & \chi' \end{pmatrix}^{-1}(x, \lambda), \quad x \in [0, 1], \quad \lambda \in \mathbb{C}. \quad (2.9)$$

Using identities (2.4), (2.5), we obtain

$$\begin{pmatrix} \varphi & \chi \\ \varphi' & \chi' \end{pmatrix}^{-1}(x, \lambda) = \begin{pmatrix} \varphi^{-1}(1, \lambda) & 0 \\ 0 & (\varphi^{-1})^*(1, \bar{\lambda}) \end{pmatrix} \begin{pmatrix} (\chi')^* & -\chi^* \\ -(\varphi')^* & \varphi^* \end{pmatrix}(x, \bar{\lambda}). \quad (2.10)$$

Therefore, $K(x, \lambda)$ satisfies the differential equation

$$K'(x, \lambda) = \begin{pmatrix} 0 & I \\ \tilde{V}(x) - \lambda & 0 \end{pmatrix} \cdot K(x, \lambda) - K(x, \lambda) \cdot \begin{pmatrix} 0 & I \\ V(x) - \lambda & 0 \end{pmatrix}, \quad x \in [0, 1]. \quad (2.11)$$

It follows from Proposition 2.3 that $K(x, \lambda)$ satisfies the initial condition

$$K(1, \lambda) = \begin{pmatrix} \tilde{\varphi}(1, \lambda) & 0 \\ \tilde{\varphi}'(1, \lambda) & I \end{pmatrix} \begin{pmatrix} \varphi^{-1}(1, \lambda) & 0 \\ -[\varphi' \varphi^{-1}](1, \lambda) & I \end{pmatrix} = \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix}.$$

Then, each function $K(x, \cdot)$, $x \in [0, 1]$, is entire as a solution of equation (2.11).

Fix some $x \in [0, 1]$. Substituting asymptotics (2.1), (2.2) into (2.9), (2.10), we obtain

$$K(x, z^2) = \begin{pmatrix} I + O(|z|^{-1}) & O(|z|^{-2}) \\ O(1) & I + O(|z|^{-1}) \end{pmatrix} \quad \text{as } |z| = \pi(n + \tfrac{1}{2}) \rightarrow \infty.$$

and $K(x, z^2) \rightarrow I_{2N}$ as $z \rightarrow i\infty$. Hence, $K(x, \lambda) = I_{2N}$ for each $(x, \lambda) \in [0, 1] \times \mathbb{C}$. In particular, this gives $\tilde{\varphi}(x, \lambda) = \varphi(x, \lambda)$ for all $(x, \lambda) \in [0, 1] \times \mathbb{C}$, i.e. $\tilde{V} = V$. $\square$

Introduce the subspaces

$$\mathcal{E}_\alpha^\# = \text{Ker } \varphi^*(1, \lambda_\alpha, V), \quad \alpha \geq 1. \quad (2.12)$$

and let $P_\alpha^\# : \mathbb{C}^N \rightarrow \mathcal{E}_\alpha^\#$ be the orthogonal projector. Using (2.5) and (2.3), we obtain $\mathcal{E}_\alpha^\#(V) = \mathcal{E}_\alpha(V^\#)$, $P_\alpha^\#(V) = P_\alpha(V^\#)$ for all $\alpha \geq 1$, where $V^\#(t) = V(1 - t)$, $t \in [0, 1]$.

Lemma 2.4. *Let $V = V^* \in L^1(0, 1)$ and $\alpha \geq 1$. The following identities are fulfilled:*

$$P_\alpha^\# \varphi'(1, \lambda_\alpha) P_\alpha = \varphi'(1, \lambda_\alpha) P_\alpha, \quad \chi'(0, \lambda_\alpha) \varphi'(1, \lambda_\alpha) P_\alpha = P_\alpha, \quad (2.13)$$

$$\text{res}_{\lambda=\lambda_\alpha} \chi^{-1}(0, \lambda_\alpha) \cdot \dot{\chi}(0, \lambda_\alpha) P_\alpha^\# = P_\alpha^\#. \quad (2.14)$$

Proof. Due to (2.4), we have $[\varphi^* \varphi'](1, \lambda_\alpha) P_\alpha = [(\varphi')^* \varphi](1, \lambda_\alpha) P_\alpha = 0$. This yields the first identity in (2.13). Let $\eta(x) = \chi(x, \lambda_\alpha) \varphi'(1, \lambda_\alpha) P_\alpha - \varphi(x, \lambda_\alpha) P_\alpha$. Then the function η satisfies the equation $-\eta'' + V\eta = \lambda_\alpha \eta$ and

$$\eta(1) = -\varphi(1, \lambda_\alpha) P_\alpha = 0, \quad \eta'(1) = \varphi'(1, \lambda_\alpha) P_\alpha - \varphi'(1, \lambda_\alpha) P_\alpha = 0.$$

Therefore, $\eta(x) = 0$ for all $x \in [0, 1]$. Using $\eta'(0) = 0$, we obtain the second identity in (2.13). Furthermore, asymptotics (2.8) yields

$$\operatorname{res}_{\lambda=\lambda_\alpha} \varphi^{-1}(1, \lambda) \cdot \dot{\varphi}(1, \lambda_\alpha) P_\alpha = P_\alpha Z_\alpha^{-1} \cdot \dot{\varphi}(1, \lambda_\alpha) P_\alpha = P_\alpha Z_\alpha^{-1} \cdot Z_\alpha P_\alpha = P_\alpha,$$

since $Z_\alpha = \dot{\varphi}(1, \lambda_\alpha) P_\alpha + \varphi(1, \lambda_\alpha) P_\alpha^\perp$. Applying this formula with the potential $V^\sharp$ instead of V and using (2.3) and $\mathcal{P}_\alpha(V^\sharp) = \mathcal{P}_\alpha^\sharp(V)$, we obtain (2.14). $\square$

Proof of Proposition 1.6. Identity (2.4) gives $m(\lambda) = m^*(\bar{\lambda})$. This implies

$$B_\alpha = - \operatorname{res}_{\lambda=\lambda_\alpha} m(\lambda) = B_\alpha^*$$

for all $\alpha \geq 1$. Due to (2.8), the function $\chi^{-1}(1, \lambda) = -(\varphi^*)^{-1}(1, \bar{\lambda})$ has a simple pole at each point $\lambda = \lambda_\alpha$. Therefore, the function $m(\lambda)$ has a simple pole at $\lambda = \lambda_\alpha$ and

$$B_\alpha = \chi'(0, \lambda_\alpha) \cdot \operatorname{res}_{\lambda=\lambda_\alpha} (\varphi^*)^{-1}(1, \bar{\lambda}) = \chi'(0, \lambda_\alpha) (Z_\alpha^*)^{-1} P_\alpha.$$

This yields $B_\alpha|_{\mathbb{C}^N \ominus \mathcal{E}_\alpha} = 0$. Recall that $G_\alpha = P_\alpha S_\alpha P_\alpha = P_\alpha [\dot{\varphi}^* \varphi'](1, \lambda_\alpha) P_\alpha$ (see (2.6)). Hence,

$$B_\alpha G_\alpha = B_\alpha [\dot{\varphi}^* \varphi'](1, \lambda_\alpha) P_\alpha = \chi'(0, \lambda_\alpha) \cdot \operatorname{res}_{\lambda=\lambda_\alpha} \chi^{-1}(0, \lambda) \cdot \dot{\chi}(0, \lambda_\alpha) \varphi'(1, \lambda_\alpha) P_\alpha,$$

where we have used the identity $\dot{\varphi}^*(1, \lambda_\alpha) = -\dot{\chi}(0, \lambda_\alpha)$. Applying Lemma 2.4, we obtain

$$\begin{aligned} B_\alpha G_\alpha &= \chi'(0, \lambda_\alpha) \cdot \operatorname{res}_{\lambda=\lambda_\alpha} \chi^{-1}(0, \lambda) \cdot \dot{\chi}(0, \lambda_\alpha) P_\alpha^\sharp \varphi'(1, \lambda_\alpha) P_\alpha \\ &= \chi'(0, \lambda_\alpha) P_\alpha^\sharp \varphi'(1, \lambda_\alpha) P_\alpha = \chi'(0, \lambda_\alpha) \varphi'(1, \lambda_\alpha) P_\alpha = P_\alpha, \quad \alpha \geq 1. \end{aligned}$$

Therefore, $B_\alpha|_{\mathcal{E}_\alpha} = g_\alpha^{-1}$ for all $\alpha \geq 1$. $\square$

3 Isospectral transforms

Let $V = V^* \in L^1(0, 1)$ and $\alpha \geq 1$. We will use the notations

$$\varphi_\alpha(x) = \varphi(x, \lambda_\alpha), \quad S_\alpha(x) = \int_0^x [\varphi_\alpha^* \varphi_\alpha](t) dt, \quad S_\alpha = S_\alpha(1).$$

Recall that the residues of the Weyl-Titchmarsh function are given by

$$B_\alpha = - \operatorname{res}_{\lambda=\lambda_\alpha} m(\lambda, V) = B_\alpha^*, \quad B_\alpha|_{\mathcal{E}_\alpha} = g_\alpha^{-1}, \quad B_\alpha|_{\mathbb{C}^N \ominus \mathcal{E}_\alpha} = 0 \quad (3.1)$$

(see Proposition 1.6). Note that $B_\alpha \geq 0$ and **the matrix B_α is uniquely determined by the pair $\{\mathcal{E}_\alpha, g_\alpha\}$ and visa versa.** In particular, $\mathcal{E}_\alpha = \mathbb{C}^N \ominus \operatorname{Ker} B_\alpha$.

Lemma 3.1. *Let $V = V^* \in L^1(0, 1)$, $\alpha \geq 1$ and let e_α be some $k_\alpha \times N$ matrix whose columns form the basis of the subspace $\mathcal{E}_\alpha$. Then*

$$B_\alpha = e_\alpha [e_\alpha^* S_\alpha e_\alpha]^{-1} e_\alpha^*. \quad (3.2)$$

Proof. If the columns of e_α form the orthonormal basis in $\mathcal{E}_\alpha$, then (3.1) and (3.2) are equivalent. Let e'_α be another $k_\alpha \times N$ matrix whose columns form the basis of $\mathcal{E}_\alpha$. Then, $e_\alpha = e'_\alpha U$ for some $k_\alpha \times k_\alpha$ matrix U such that $\det U \neq 0$. The simple calculation shows that

$$B_\alpha(V) = e'_\alpha U [U^* (e'_\alpha)^* S_\alpha (1) e'_\alpha U]^{-1} U^* (e'_\alpha)^* = e'_\alpha [(e'_\alpha)^* S_\alpha (1) e'_\alpha]^{-1} (e'_\alpha)^*.$$

Hence, (3.2) doesn't depend on the choice of e_α . $\square$

Introduce the matrices

$$D_\alpha = S_\alpha^{-1} - B_\alpha, \quad \alpha \geq 1.$$

Lemma 3.2. *Let $V = V^* \in L^1(0, 1)$, $\alpha \geq 1$ and the subspace $\mathcal{F}_\alpha \subset \mathbb{C}^N$ be given by (1.4). Then $D_\alpha = D_\alpha^* \geq 0$, $\mathcal{F}_\alpha = \mathbb{C}^N \ominus \text{Ker } D_\alpha$ and $\dim \mathcal{F}_\alpha = N - k_\alpha$. Moreover, $\mathcal{F}_\alpha \cap \mathcal{E}_\alpha = \{0\}$.*

Proof. Without loss of generality, we can assume that

$$e_\alpha = \begin{pmatrix} I_{k_\alpha} \\ 0 \end{pmatrix}; \quad S_\alpha = \begin{pmatrix} s & p \\ p^* & q \end{pmatrix}, \quad B_\alpha = e_\alpha (e_\alpha^* S_\alpha e_\alpha)^{-1} e_\alpha^* = \begin{pmatrix} s^{-1} & 0 \\ 0 & 0 \end{pmatrix},$$

where $s = s^*$ is a $k_\alpha \times k_\alpha$ matrix, p is $(N - k_\alpha) \times k_\alpha$ and $q = q^*$ is $(N - k_\alpha) \times (N - k_\alpha)$. Note that $S_\alpha > 0$ yields $s > 0$ and $q > 0$. Due to the Frobenius formula for the inverse matrix (see [G], Ch. 2.5), we have

$$S_\alpha^{-1} = \begin{pmatrix} (s - pq^{-1}p^*)^{-1} & -s^{-1}p(q - p^*s^{-1}p)^{-1} \\ -q^{-1}p^*(s - pq^{-1}p^*)^{-1} & (q - p^*s^{-1}p)^{-1} \end{pmatrix}.$$

Note that $q^{-1}p^*(s - pq^{-1}p^*)^{-1} = (q - p^*s^{-1}p)^{-1}p^*s^{-1}$, since $S_\alpha^{-1} = (S_\alpha^{-1})^*$. Together with the identity $(s - pq^{-1}p^*)^{-1} - s^{-1} = s^{-1}pq^{-1}p^*(s - pq^{-1}p^*)^{-1}$, this yields

$$\begin{aligned} D_\alpha = S_\alpha^{-1} - B_\alpha &= \begin{pmatrix} s^{-1}pq^{-1}p^*(s - pq^{-1}p^*)^{-1} & -s^{-1}p(q - p^*s^{-1}p)^{-1} \\ -q^{-1}p^*(s - pq^{-1}p^*)^{-1} & (q - p^*s^{-1}p)^{-1} \end{pmatrix} \\ &= \begin{pmatrix} -s^{-1}p \\ I_{N-k_\alpha} \end{pmatrix} (q - p^*s^{-1}p)^{-1} \begin{pmatrix} -p^*s^{-1} & I_{N-k_\alpha} \end{pmatrix}. \end{aligned}$$

This implies $D_\alpha \geq 0$, $\text{rank } D_\alpha = N - k_\alpha$ and $\dim \text{Ker } D_\alpha = k_\alpha$. Moreover, the identity

$$\begin{pmatrix} -p^*s^{-1} & I_{N-k_\alpha} \end{pmatrix} \begin{pmatrix} s \\ p^* \end{pmatrix} = 0$$

yields $S_\alpha(\mathcal{E}_\alpha) \subset \text{Ker } D_\alpha$. Recall that $S_\alpha = S_\alpha^* > 0$. Using $\dim \mathcal{E}_\alpha = k_\alpha = \dim \text{Ker } D_\alpha$, we deduce that $S_\alpha(\mathcal{E}_\alpha) = \text{Ker } D_\alpha$ and so $\mathcal{F}_\alpha = \mathbb{C}^N \ominus \text{Ker } D_\alpha$. Since $D_\alpha + B_\alpha > 0$, we have $\text{Ker } D_\alpha \cap \text{Ker } B_\alpha = (\mathbb{C}^N \ominus \mathcal{F}_\alpha) \cap (\mathbb{C}^N \ominus \mathcal{E}_\alpha) = \{0\}$. Together with $\dim \mathcal{F}_\alpha + \dim \mathcal{E}_\alpha = N$, this implies $\mathcal{F}_\alpha \cap \mathcal{E}_\alpha = \{0\}$. $\square$

Corollary 3.3. *Let $V = V^* \in L^1(0, 1)$. Then $\mathcal{F}_\alpha = \mathbb{C}^N \ominus [\dot{\varphi}^*(1, \lambda_\alpha)](\mathcal{E}_\alpha^\#)$ for all $\alpha \geq 1$, where the subspace $\mathcal{E}_\alpha^\# \subset \mathbb{C}^N$ is given by (2.12).*

Proof. Recall that $\mathcal{F}_\alpha = \mathbb{C}^N \ominus S_\alpha(\mathcal{E}_\alpha)$ and $S_\alpha = \int_0^1 [\varphi^* \varphi](t, \lambda_\alpha) dt = [\dot{\varphi}^* \varphi' - (\dot{\varphi}')^* \varphi](1, \lambda_\alpha)$. Since $\varphi(1, \lambda_\alpha)|_{\mathcal{E}_\alpha} = 0$, we obtain

$$S_\alpha(\mathcal{E}_\alpha) = [\dot{\varphi}^*(1, \lambda_\alpha) \varphi'(1, \lambda_\alpha)](\mathcal{E}_\alpha).$$

Using (2.13), we deduce that $[\varphi'(1, \lambda_\alpha)](\mathcal{E}_\alpha) = \mathcal{E}_\alpha^\#$. Hence, $S_\alpha(\mathcal{E}_\alpha) = [\dot{\varphi}^*(1, \lambda_\alpha)](\mathcal{S}_\alpha^\#)$. $\square$

The following Theorem gives the explicit formula for the isospectral transform of the potential V , changing only the matrix $B_\alpha(V)$.

Theorem 3.4. *Let the potential $V = V^* \in L^1(0, 1)$ and the matrix $B = B^*$ be such that*

$$B \geq 0, \quad \text{rank } B = k_\alpha \quad \text{and} \quad \mathcal{E}^{(B)} \cap \mathcal{F}_\alpha = \{0\}, \quad \text{where} \quad \mathcal{E}^{(B)} = \mathbb{C}^N \ominus \text{Ker } B, \quad (3.3)$$

for some $\alpha \geq 1$. Denote

$$\tilde{V}(x) = V(x) - 2[\varphi_\alpha K \varphi_\alpha^*]'(x) = \tilde{V}^*(x), \quad x \in [0, 1], \quad (3.4)$$

where

$$K(x) = A(I + S_\alpha(x)A)^{-1} = K^*(x) \quad \text{and} \quad A = B - B_\alpha. \quad (3.5)$$

Then $(\tilde{V} - V)' \in L^1(0, 1)$, $\tilde{B} = B_\beta$ for all $\beta \neq \alpha$, and

$$\tilde{B}_\alpha = B, \quad \tilde{\mathcal{E}}_\alpha = \mathcal{E}^{(B)}, \quad \tilde{\mathcal{F}}_\alpha = \mathcal{F}_\alpha.$$

In order to prove Theorem 3.4, we need two preliminary Lemmas.

Lemma 3.5. *Let $V = V^* \in L^1(0, 1)$, $\alpha \geq 1$ and $B = B^*$ be such that (3.3) hold true. Then*

(i) $\det(I + S_\alpha(x)A) \neq 0$ for all $x \in [0, 1]$, where $A = B - B_\alpha$.

(ii) the identity $[\varphi_\alpha K \varphi_\alpha^*](1) = 0$ is fulfilled, where K is given by (3.5).

Proof. (i) Note that $I + S_\alpha(x)A = S_\alpha(x)(S_\alpha^{-1}(x) + A)$. For all $x \in [0, 1]$ we have

$$S_\alpha^{-1}(x) + A > S_\alpha^{-1}(1) + A = (S_\alpha^{-1}(1) - B_\alpha) + (B_\alpha + A) = D_\alpha + B \geq 0,$$

since $D_\alpha \geq 0$ and $B \geq 0$. Lemma 3.2 and (3.3) give $\text{rank } D_\alpha = N - k_\alpha$, $\text{rank } B = k_\alpha$ and $(\mathbb{C}^N \ominus \text{Ker } D_\alpha) \cap (\mathbb{C}^N \ominus \text{Ker } B) = \{0\}$. Hence, $\text{Ker } D_\alpha \cap \text{Ker } B = \{0\}$ and $D_\alpha + B > 0$.

(ii) Let the columns of the matrix e_α form some orthonormal basis in E_α . Without loss of generality, we can assume that

$$e_\alpha = \begin{pmatrix} I_{k_\alpha} \\ 0 \end{pmatrix}; \quad S_\alpha(1) = \begin{pmatrix} s & p \\ p^* & q \end{pmatrix}, \quad B = \begin{pmatrix} b & c \\ c^* & d \end{pmatrix},$$

where $s = s^*$ and $b = b^*$ are $k_\alpha \times k_\alpha$ matrices, p and c are $(N - k_\alpha) \times k_\alpha$ matrices, $q = q^*$ and $d = d^*$ are $(N - k_\alpha) \times (N - k_\alpha)$ matrices. Then,

$$B_\alpha = e_\alpha(e_\alpha^* S_\alpha(1) e_\alpha)^{-1} e_\alpha^* = \begin{pmatrix} s^{-1} & 0 \\ 0 & 0 \end{pmatrix}, \quad A = \begin{pmatrix} b - s^{-1} & c \\ c^* & d \end{pmatrix}.$$

Firstly, let $b > 0$. In this case, the assumption $\text{rank } B = k_\alpha$ gives $d = c^*b^{-1}c$. Note that

$$I + S_\alpha(1)A = \begin{pmatrix} sb + pc^* & sc + pd \\ p^*(b - s^{-1}) + qc^* & I_{N-k_\alpha} + p^*c + qd \end{pmatrix}$$

and

$$\begin{pmatrix} sb + pc^* & sc + pd \end{pmatrix} = (s + pc^*b^{-1}) \cdot \begin{pmatrix} b & c \end{pmatrix}. \quad (3.6)$$

Moreover,

$$\begin{pmatrix} c^* & d \end{pmatrix} = c^*b^{-1} \cdot \begin{pmatrix} b & c \end{pmatrix}. \quad (3.7)$$

Due to (3.6), (3.7), for each index $j \in [k_\alpha + 1, N]$, the first k_α rows of the matrix $I + S_\alpha(1)A$ and the j -th row of A are linearly dependent. Using the Cramer formula¹, we deduce that the matrix $K(1) = A(I + S_\alpha(1)A)^{-1}$ has the form

$$K(1) = \begin{pmatrix} \cdots & \cdots \\ c^* & d \end{pmatrix} \begin{pmatrix} sb + pc^* & sc + pd \\ \cdots & \cdots \end{pmatrix}^{-1} = \begin{pmatrix} \cdots & \cdots \\ \cdots & 0 \end{pmatrix}.$$

Recall that $\varphi_\alpha(1)e_\alpha = 0$. Hence, $\varphi_\alpha(1) = \begin{pmatrix} 0 & \cdots \\ 0 & \cdots \end{pmatrix}$. This yields $[\varphi_\alpha^* K \varphi_\alpha](1) = 0$.

Secondly, let $\det b = 0$. Since $K(1)$ is a continuous function of B , we deduce that it has the same form as before and $[\varphi_\alpha^* K \varphi_\alpha](1) = 0$. $\square$

Introduce the matrices

$$S(x, \lambda) = \int_0^x [\varphi^* \varphi](t, \lambda) dt, \quad \tilde{S}(x, \lambda) = \int_0^x [\tilde{\varphi}^* \tilde{\varphi}](t, \lambda) dt.$$

Lemma 3.6. *Let the potential $V = V^* \in L^1(0, 1)$, the number $\alpha \geq 1$ and the matrix $B = B^*$ be such that conditions (3.3) are fulfilled. Then*

$$\tilde{\varphi}(x, \lambda) = \varphi(x, \lambda) - [\varphi_\alpha K T](x, \lambda) \quad \text{and} \quad \tilde{S}(x, \lambda) = S(x, \lambda) - [T^* K T](x, \lambda),$$

where

$$T(x, \lambda) = \int_0^x \varphi_\alpha^*(t) \varphi(t, \lambda) dt$$

and the potential $\tilde{V}$ is given by (3.4).

¹Let $X = [x_{j,m}]_{j,m=1}^N$, $Y = [y_{j,m}]_{j,m=1}^N$ be $N \times N$ matrices and let $\det Y \neq 0$, $Z = XY^{-1} = [z_{j,m}]_{j,m=1}^N$. Then

$$z_{j,m} = \frac{1}{\det Y} \begin{vmatrix} y_{1,1} & y_{1,2} & \cdots & y_{1,N-1} & y_{1,N} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ y_{m-1,1} & y_{m-1,2} & \cdots & y_{m-1,N-1} & y_{m-1,N} \\ x_{j,1} & x_{j,2} & \cdots & x_{j,N-1} & x_{j,N} \\ y_{m+1,1} & y_{m+1,2} & \cdots & y_{m+1,N-1} & y_{m+1,N} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ y_{N,1} & y_{N,2} & \cdots & y_{N,N-1} & y_{N,N} \end{vmatrix} \quad \text{for all } j, m \in [1, N] \quad (\text{see [G], Ch. 1.3}).$$

Proof. Let $\eta = \varphi - \varphi_\alpha KT$ (we omit x and λ for short). Using the identity $K' = -K\varphi_\alpha^* \varphi_\alpha K$, we obtain $(KT)' = K\varphi_\alpha^* \eta$. Therefore,

$$\eta' = \varphi' - \varphi'_\alpha KT - \varphi_\alpha K\varphi_\alpha^* \eta,$$

$$\eta'' = \varphi'' - \varphi''_\alpha KT - \varphi'_\alpha K\varphi_\alpha^* \eta - (\varphi_\alpha K\varphi_\alpha^*)' \eta - \varphi_\alpha K\varphi_\alpha^* (\varphi' - \varphi'_\alpha KT - \varphi_\alpha K\varphi_\alpha^* \eta).$$

Recall that $\varphi_\alpha^* \varphi'_\alpha = (\varphi'_\alpha)^* \varphi_\alpha$ (see (2.4)). Hence,

$$\eta'' = \varphi'' - \varphi''_\alpha KT - 2(\varphi_\alpha K\varphi_\alpha^*)' \eta + \varphi_\alpha K((\varphi'_\alpha)^* \varphi - \varphi_\alpha^* \varphi').$$

Note that $\varphi'' = (V - \lambda)\varphi$, $\varphi''_\alpha = (V - \lambda_\alpha)\varphi_\alpha$ and $(\varphi'_\alpha)^* \varphi - \varphi_\alpha^* \varphi' = (\lambda - \lambda_\alpha)T$. This gives

$$\eta'' = (V - \lambda)\eta - 2(\varphi_\alpha K\varphi_\alpha^*)' \eta = (\tilde{V} - \lambda)\eta.$$

Since $\eta(0) = 0$ and $\eta'(0) = \varphi'(0) = I$, we deduce $\eta = \tilde{\varphi}$. Furthermore,

$$\eta^* \eta = \varphi^* \varphi - \varphi^* \varphi_\alpha KT + T^* K\varphi_\alpha^* \varphi_\alpha KT - T^* K\varphi_\alpha^* \varphi = \varphi^* \varphi - [T^* KT]'.$$

This yields $\tilde{S} = S - T^* KT$. □

Proof of Theorem 3.4. Note that $(\varphi_\alpha K\varphi_\alpha^*)'' \in L^1(0, 1)$, since $\varphi''_\alpha, K'' \in L^1(0, 1)$. Fix some $\beta \neq \alpha$. Let $\varphi_\beta(x) = \varphi(x, \lambda_\beta, V)$, $\tilde{\varphi}_\beta(x) = \varphi(x, \lambda_\beta, \tilde{V})$ and so on. Using Lemma 3.6 and the identity

$$T_\beta(1) = \int_0^1 [\varphi_\alpha^* \varphi_\beta](t) dt = \frac{[\varphi_\alpha^* \varphi'_\beta - (\varphi'_\alpha)^* \varphi_\beta](1)}{\lambda_\alpha - \lambda_\beta}, \quad (3.8)$$

we obtain

$$\tilde{\varphi}_\beta(1) = \varphi_\beta(1) - \frac{[\varphi_\alpha K(\varphi_\alpha^* \varphi'_\beta - (\varphi'_\alpha)^* \varphi_\beta)](1)}{\lambda_\alpha - \lambda_\beta} = \left(I + \frac{[\varphi_\alpha K(\varphi'_\alpha)^*](1)}{\lambda_\alpha - \lambda_\beta} \right) \varphi_\beta(1), \quad (3.9)$$

where we have used $[\varphi_\alpha K\varphi_\alpha^*](1) = 0$ (see Lemma 3.5 (ii)). Therefore, λ_β is a root of the (scalar) entire function $\tilde{w}(\lambda) = \det \tilde{\varphi}(1, \lambda)$ of the multiplicity at least k_β . Furthermore,

$$\tilde{\varphi}_\alpha(1) = \varphi_\alpha(1) - [\varphi_\alpha K S_\alpha](1) = \varphi_\alpha(1) (I - [K S_\alpha](1)). \quad (3.10)$$

Hence, λ_α is a root of $\tilde{w}(\lambda)$ of the multiplicity at least k_α . Using Lemma 2.2 (ii), we deduce that $[\tilde{w}w^{-1}](\lambda)$ is an entire function, where $w(\lambda) = \det \varphi(1, \lambda)$. Note that $[\tilde{w}w^{-1}](z^2) = 1 + O(|z|^{-1})$ as $|z| = \pi^2(n + \frac{1}{2})^2 \rightarrow \infty$ (see (2.1)). Hence, $\tilde{w}(\lambda) = w(\lambda)$, $\lambda \in \mathbb{C}$. In other words, each λ_β , $\beta \geq 1$, is an eigenvalue of the operator $\tilde{H}\psi = -\psi'' + \tilde{V}\psi$ of the multiplicity k_β and there are no other eigenvalues.

Let $\beta \neq \alpha$. Identity (3.9) gives $\tilde{\mathcal{E}}_\beta \supset \mathcal{E}_\beta$. Since $\dim \tilde{\mathcal{E}}_\beta = k_\beta = \dim \mathcal{E}_\beta$, we obtain $\tilde{\mathcal{E}}_\beta = \mathcal{E}_\beta$. Let e_β be some $k_\beta \times N$ matrix whose columns form the basis of E_β . Recall that $\tilde{B}_\beta = e_\beta(e_\beta^* \tilde{S}_\beta e_\beta)^{-1} e_\beta^*$. Using Lemma 3.6 and (3.8), we obtain

$$\tilde{S}_\beta = S_\beta - \frac{[(\varphi'_\beta)^* \varphi_\alpha - \varphi_\beta^* \varphi'_\alpha] K(\varphi_\alpha^* \varphi'_\beta - (\varphi'_\alpha)^* \varphi_\beta)](1)}{(\lambda_\alpha - \lambda_\beta)^2}.$$

Note that $\varphi_\beta(1)e_\beta = 0$, since the columns of e_β belong to $\mathcal{E}_\beta = \text{Ker } \varphi_\beta(1)$. Due to Lemma 3.5, $[\varphi_\alpha^* K \varphi_\alpha](1) = 0$. Therefore, $e_\beta^* \tilde{S}_\beta e_\beta = e_\beta^* S_\beta e_\beta$ and

$$\tilde{B}_\beta = e_\beta(e_\beta^* \tilde{S}_\beta e_\beta)^{-1} e_\beta^* = e_\beta(e_\beta^* S_\beta e_\beta)^{-1} e_\beta^* = B_\beta, \quad \beta \neq \alpha.$$

We will show that $\tilde{B}_\alpha = B$. Let the columns of the matrix e_α form some orthonormal basis in E_α . Without loss of generality, we can assume that

$$e_\alpha = \begin{pmatrix} I_{k_\alpha} \\ 0 \end{pmatrix}; \quad S_\alpha = \begin{pmatrix} s & p \\ p^* & q \end{pmatrix}, \quad B = \begin{pmatrix} b & c \\ c^* & d \end{pmatrix},$$

where $s = s^*$ and $b = b^*$ are $k_\alpha \times k_\alpha$ matrices, p and c are $(N - k_\alpha) \times k_\alpha$ matrices, $q = q^*$ and $d = d^*$ are $(N - k_\alpha) \times (N - k_\alpha)$ matrices. Note that

$$B_\alpha = e_\alpha(e_\alpha^* S e_\alpha)^{-1} e_\alpha^* = \begin{pmatrix} s^{-1} & 0 \\ 0 & 0 \end{pmatrix}, \quad A = \begin{pmatrix} b - s^{-1} & c \\ c^* & d \end{pmatrix}.$$

Firstly, let $b > 0$. In this case, $\text{rank } B = k_\alpha$ implies $d = c^* b^{-1} c$. It follows from identity (3.10) that the columns of the matrix

$$\tilde{e}_\alpha = (I - K S_\alpha)^{-1} e_\alpha = (I - A(S_\alpha^{-1} + A)^{-1})^{-1} e_\alpha = (I + A S_\alpha) e_\alpha$$

form a basis in $\tilde{E}_\alpha$ (recall that $\det(I + A S_\alpha) = \det(I + S_\alpha A) \neq 0$ due to Lemma 3.5 (i)). We have

$$\tilde{e}_\alpha = \begin{pmatrix} I_{k_\alpha} + (b - s^{-1})s + c p^* \\ c^* s + d p^* \end{pmatrix} = \begin{pmatrix} b s + c p^* \\ c^* s + c^* b^{-1} c p^* \end{pmatrix} = \begin{pmatrix} I_{k_\alpha} \\ c^* b^{-1} \end{pmatrix} (b s + c p^*).$$

Furthermore, Lemma 3.6 gives

$$\tilde{S}_\alpha = S_\alpha - S_\alpha^* K S_\alpha = S_\alpha (I - K S_\alpha) = S_\alpha (I + A S_\alpha)^{-1}.$$

Hence,

$$\tilde{e}_\alpha^* \tilde{S}_\alpha \tilde{e}_\alpha = \tilde{e}_\alpha^* S_\alpha e_\alpha = (s b + p c^*) \begin{pmatrix} I_{k_\alpha} & b^{-1} c \end{pmatrix} \begin{pmatrix} s \\ p^* \end{pmatrix} = (s b + p c^*) b^{-1} (b s + c p^*).$$

We obtain

$$\tilde{B}_\alpha = \tilde{e}_\alpha (\tilde{e}_\alpha^* \tilde{S}_\alpha \tilde{e}_\alpha)^{-1} \tilde{e}_\alpha^* = \begin{pmatrix} I_{k_\alpha} \\ c^* b^{-1} \end{pmatrix} \cdot b \cdot \begin{pmatrix} I_{k_\alpha} & b^{-1} c \end{pmatrix} = \begin{pmatrix} b & c \\ c^* & c^* b^{-1} c \end{pmatrix} = B.$$

Secondly, let $\det b = 0$. Note that $\tilde{e}_\alpha$ and $\tilde{S}_\alpha$ are continuous functions of B . Therefore, $\tilde{B}_\alpha = B$ due to the arguments given above and the continuity of $\tilde{B}_\alpha$ as a function of B . Since $\tilde{B}_\alpha = B$, we obtain $\tilde{\mathcal{E}}_\alpha = \mathbb{C}^N \ominus \text{Ker } B = \mathcal{E}^{(B)}$. Note that

$$\tilde{S}_\alpha^{-1} = (S_\alpha (I + A S_\alpha)^{-1})^{-1} = S_\alpha^{-1} + A.$$

This implies

$$\tilde{D}_\alpha = \tilde{S}_\alpha^{-1} - \tilde{B}_\alpha = S_\alpha^{-1} + A - B = S_\alpha^{-1} - B_\alpha = D_\alpha.$$

In particular, we have $\tilde{\mathcal{F}}_\alpha = \mathbb{C}^N \ominus \text{Ker } \tilde{D}_\alpha = \mathbb{C}^N \ominus \text{Ker } D_\alpha = \mathcal{F}_\alpha$. □

Proof of Theorem 1.3. Due to Theorem 1.2, the mapping Φ_α is one-to-one. We prove that Φ_α is onto. Let the subspace $\mathcal{E} \subset \mathbb{C}^N$ be such that $\dim \mathcal{E} = k_\alpha$, $\mathcal{E} \cap \mathcal{F}_\alpha = \{0\}$ and let $g = g^* > 0$ be the operator in $\mathcal{E}$. We define the matrix $B = B^* \geq 0$ by

$$B|_{\mathcal{E}} = g^{-1}, \quad B|_{\mathbb{C}^N \ominus \mathcal{E}} = 0.$$

Since $\mathcal{E}^{(B)} = \mathbb{C}^N \ominus \text{Ker } B = \mathcal{E}$, conditions (3.3) are fulfilled. Let $\tilde{V}$ be given by formula (3.4). It follows from Theorem 3.4 that $\tilde{V} \in \text{Iso}(V)$, $\tilde{B}_\beta = B_\beta$ for all $\beta \neq \alpha$ and $\tilde{B}_\alpha = B$. In view of definition (3.1), this yields $\tilde{\mathcal{E}}_\beta = \mathcal{E}_\beta$, $\tilde{g}_\beta = g_\beta$ for all $\beta \neq \alpha$ and $\tilde{\mathcal{E}}_\alpha = \mathcal{E}$, $\tilde{g}_\alpha = g$. $\square$

Proof of Proposition 1.4. Since $\dim \mathcal{F}_\alpha = \dim \tilde{\mathcal{F}}_\alpha = N - k_\alpha$, we can fix some matrix $B = B^* \geq 0$ such that $\text{rank } B = k_\alpha$, $\mathcal{E}^{(B)} \cap \mathcal{F}_\alpha = \{0\}$ and $\mathcal{E}^{(B)} \cap \tilde{\mathcal{F}}_\alpha = \{0\}$, where $\mathcal{E}^{(B)} = \mathbb{C}^N \ominus \text{Ker } B$. Using Theorem 3.4, we construct potentials $V_B, \tilde{V}_B \in \text{Iso}(V)$ such that

$$B_\beta(V_B) = B_\beta, \quad B_\beta(\tilde{V}_B) = \tilde{B}_\beta, \quad \beta \neq \alpha, \quad B_\alpha(V_B) = B = B_\alpha(\tilde{V}_B).$$

Note that $\mathcal{F}_\alpha(V_B) = \mathcal{F}_\alpha$ and $\mathcal{F}_\alpha(\tilde{V}_B) = \tilde{\mathcal{F}}_\alpha$. It is clear that

$$\mathcal{E}_\beta(V_B) = \mathcal{E}_\beta = \tilde{\mathcal{E}}_\beta = \mathcal{E}_\beta(\tilde{V}_B), \quad \beta \neq \alpha, \quad \text{and} \quad \mathcal{E}_\alpha(V_B) = \mathcal{E}^{(B)} = \mathcal{E}_\alpha(\tilde{V}_B).$$

Due to Proposition 2.3 (i), we have $\varphi(1, \lambda, V_B) = \varphi(1, \lambda, \tilde{V}_B)$ for all $\lambda \in \mathbb{C}$. In particular, $\dot{\varphi}^*(1, \lambda_\alpha, V_B) = \dot{\varphi}^*(1, \lambda_\alpha, \tilde{V}_B)$ and $\mathcal{E}_\alpha^\#(V_B) = \text{Ker } \varphi^*(1, \lambda_\alpha, V_B) = \text{Ker } \varphi^*(1, \lambda_\alpha, \tilde{V}_B) = \mathcal{E}_\alpha^\#(\tilde{V}_B)$. Corollary 3.3 implies $\mathcal{F}_\alpha(V_B) = \mathcal{F}_\alpha(\tilde{V}_B)$. Therefore, $\mathcal{F}_\alpha = \tilde{\mathcal{F}}_\alpha$. Together with Lemma 3.2 this gives $\tilde{\mathcal{E}}_\alpha \cap \mathcal{F}_\alpha = \tilde{\mathcal{E}}_\alpha \cap \tilde{\mathcal{F}}_\alpha = \{0\}$. $\square$

Proof of Proposition 1.5. Let $\mathcal{E}_1 \cap \mathcal{E}_2 \neq \{0\}$. Due to $\dim \mathcal{E}_1 = \dim \mathcal{E}_2 = 1$, this is equivalent to $\mathcal{E}_1 = \mathcal{E}_2$. Fix some vector $h \in \mathcal{E}_1 = \mathcal{E}_2$ such that $\|h\| = 1$. Let

$$w(\lambda) = \det \varphi(1, \lambda), \quad u(\lambda) = [h^* \varphi(1, \lambda) h]^2, \quad \lambda \in \mathbb{C}.$$

It follows from (2.1) that both functions w, u have the same asymptotics

$$w(z^2), u(z^2) = \frac{\sin^2 z}{z^2} \cdot (1 + O(|z|^{-1})) \quad \text{as } |z| = \pi(n + \frac{1}{2}) \rightarrow \infty.$$

Therefore, by Rouché's Theorem, $w(\lambda)$ and $u(\lambda)$ have the same number of roots counted with multiplicity in the disc $\{|\lambda| < \pi^2(N + \frac{1}{2})^2\}$, if N is sufficiently large. Due to Lemma 2.2 (ii), $w(\lambda)$ has simple roots at the points λ_1, λ_2 and $w(\lambda)$ has double roots at the points λ_α , $\alpha \geq 3$. Moreover, $w(\lambda)$ has no other roots. On the other hand, each point λ_α , $\alpha \geq 1$, is a double root of $u(\lambda)$. This is a contradiction. Hence, $\mathcal{E}_1 \cap \mathcal{E}_2 = \{0\}$.

Suppose that $\mathcal{F}_1 \neq \mathcal{E}_2$, i.e. $\mathcal{F}_1 \cap \mathcal{E}_2 = \{0\}$. Then, using Theorem 1.3, we can construct the potential $\tilde{V} \in \text{Iso}(V)$ such that $\tilde{\mathcal{E}}_1 = \mathcal{E}_2 = \tilde{\mathcal{E}}_2$. Due to the arguments given above, this is impossible. Therefore, $\mathcal{F}_1 = \mathcal{E}_2$. The proof of $\mathcal{F}_2 = \mathcal{E}_1$ is similar. $\square$

Acknowledgments. Dmitry Chelkak was partly supported by grants VNP Minobrazovaniya 3.1–4733 and "Vedushie nauchnye shkoly". Evgeny Korotyaev was partly supported by DFG project BR691/23-1. Some part of this paper was written at the Mittag-Leffler Institute, Stockholm. The authors are grateful to the Institute for the hospitality. The authors would like to thank Markus Klein for useful discussions.

References

- [Ca] Carlson R.: An inverse problem for the matrix Schrödinger equation. J. Math. Anal. Appl. 267 (2002), no. 2, 564–575.
- [CK] Chelkak D., Korotyaev E.: Spectral estimates for Schrödinger operators with periodic matrix potentials on the real line. Preprint Institut Mittag-Leffler No. 06, 2005/2006 fall. To be published in International Mathematics Research Notices.
- [Ch] Chern Hua-Huai: On the construction of isospectral vectorial Sturm-Liouville differential equation. Preprint 1998.
- [ChSh] Chern Hua-Huai; Shen Chao-Liang: On the n -dimensional Ambarzumyan's theorem. Inverse Problems **13**(1), 15–18 (1997).
- [CHGL] Clark S., Holden H., Gesztesy, F., Levitan, B.: Borg-type theorem for matrix-valued Schrödinger operators. J. Diff. Eqs. **167**, 181–210 (2000).
- [G] Gantmacher F. R.: *The theory of matrices. Vol. 1.* Translated from the Russian by K. A. Hirsch. Reprint of the 1959 translation. AMS Chelsea Publishing, Providence, RI, 1998.
- [JL1] Jodeit M. Jr., Levitan B. M.: Isospectral vector-valued Sturm-Liouville problems. Lett. Math. Phys. **43**(2), 117–122 (1998).
- [JL2] Jodeit M. Jr., Levitan B. M.: A characterization of some even vector-valued Sturm-Liouville problems. Mat. Fiz. Anal. Geom. **5**(3-4), 166–181 (1998).
- [L] Levitan B.: *Inverse Sturm-Liouville problems.* Utrecht: VNU Science Press, 1987.
- [Mal] Malamud M. M.: Uniqueness of the matrix Sturm-Liouville equation given a part of the monodromy matrix, and Borg type results. *Sturm-Liouville Theory: Past and Present*, 237–270. Birkhauser, Basel, 2005.
- [M] Marchenko V.: *Sturm-Liouville operator and applications.* Basel: Birkhäuser, 1986.
- [PT] Pöschel P., Trubowitz E.: *Inverse spectral theory.* Academic Press, Boston, 1987.
- [Sh] Shen Chao-Liang: Some inverse spectral problems for vectorial Sturm-Liouville equations. Inverse Problems **17**(5), 1253–1294 (2001).
- [SP] Samsonov B. F.; Pecheritsin A. A.: Chains of Darboux transformations for the matrix Schrödinger equation. J. Phys. A **37**(1), 239–250 (2004).
- [Yu] Yurko V.A.: Inverse problems for matrix Sturm-Liouville operators. Russian J. Math. Phys. **13**(1) (2006).